arXiv:1305.0164v1 [math.AC] 1 May 2013

REGULAR SEQUENCES AND *ZD*-MODULES

SH. PAYROVI AND M. LOTFI PARSA

ABSTRACT. Let R be a Noetherian ring, I an ideal of R and M a ZD-module. Let S be a Melkersson subcategory with respect to I such that $M/IM \notin S$. We show that all maximal S-sequences on M in I, have equal length. If this common length is denoted by $S-\operatorname{depth}_I(M)$, then $S-\operatorname{depth}_I(M) = \inf\{i : H^i_I(M) \notin S\}$.

1. INTRODUCTION

Throughout this paper, R is a commutative Noetherian ring with non-zero identity, I an ideal of R, M an R-module and t an integer. For notations and terminologies not given in this paper, the reader is referred to [4] and [5] if necessary.

As a generalization of finitely generated modules, Evans [7] introduced ZD-modules. An R-module M is said to be zero-divisor module (ZD-module), if $Z_R(M)$ is a finite union of prime ideals in $\operatorname{Ass}_R M$. According to [6, Example 2.2], the class of ZD-modules is much larger than that of finitely generated modules.

Let S be a Serre subcategory of the category of R-modules. The notion of S-sequences on M, is defined by Aghapournahr and Melkersson [1]. An element a of R is called S-regular on M, if $0 :_M a \in S$. A sequence $a_1, \ldots, a_t$ is an S-sequence on M, if a_i is S-regular on $M/(a_1, \ldots, a_{i-1})M$ for $i = 1, \ldots, t$. Suppose that S is a Melkersson subcategory with respect to I, M is a finitely generated R-module and $M/IM \notin S$. In [1, Lemma 2.14(b)], it is proved that the length of all maximal S-sequences on M in I, is equal to $\inf\{i : \operatorname{Ext}^i_R(R/I, M) \notin S\}$. If this common length is denoted by $S-\operatorname{depth}_I(M)$, then $S-\operatorname{depth}_I(M) = \inf\{i : H^i_I(M) \notin S\}$; see [1, Theorem 2.18(a)].

In this paper, we generalize these results to the ZD-modules.

2. MAIN RESULTS

Recall that R is a Noetherian ring, I is an ideal of R and M is an R-module.

2010 Mathematics Subject Classification. 13C15, 13D45.

Key words and Phrases. Local cohomology, Regular sequence, Serre subcategory, ZD-Module.

Definition 2.1. A full subcategory of the category of R-modules is said to be Serre subcategory, if it is closed under taking submodules, quotients and extensions. A Serre subcategory S is said to be Melkersson subcategory with respect to I, if for any I-torsion R-module M, $0 :_M I \in S$ implies that $M \in S$. A Serre subcategory is called Melkersson subcategory, when it is a Melkersson subcategory with respect to all ideals of R.

It is proved that if a Serre subcategory is closed under taking injective hulls, then it is a Melkersson subcategory; see [1, Lemma 2.2].

Example 2.2. [1, Example 2.4] The following classes of modules are Serre subcategories closed under taking injective hulls and hence are Melkersson subcategories.

(a) The class of zero modules.
(b) The class of Artinian R-modules.
(c) The class of R-modules with finite support.
(d) The class of all R-modules M with $\dim_R M \leq k$, where k is a non-negative integer. When $k = 0$, we get the class of semiartinian modules, i.e. the modules M with $\mathrm{Supp}_R M \subseteq \mathrm{Max}(R)$.

Example 2.3. [1, Example 2.5] The class of I-cofinite Artinian modules is a Melkersson subcategory with respect to I, but is not closed under taking injective hulls.

Aghapournahr and Melkersson [1] introduced the notion of S-sequences on M as a generalization of regular sequences.

Definition 2.4. An element a of R is called S-regular on M, if $0 :_M a \in S$. A sequence $a_1, \ldots, a_t$ is an S-sequence on M, if a_i is S-regular on $M/(a_1, \ldots, a_{i-1})M$ for $i = 1, \ldots, t$. The S-sequence $a_1, \ldots, a_t$ is said to be maximal S-sequence on M, if $a_1, \ldots, a_t, y$ is not an S-sequence on M for any $y \in R$.

Example 2.5. [1, Example 2.8] Let S be as in Example 2.2. S-sequences on M are just:

(a) Poor M-sequences, [4, Definition 6.2.1].
(b) Filter-regular sequences which have been defined in the local case for finitely generated modules in [12].

(c) Generalized regular sequences which have been defined in the local case for finitely generated modules in [11, Definition 2.1].

(d) M-sequences in dimension $> k$ which have been defined for finitely generated modules in [3, Definition 2.1].

Lemma 2.6. *If $M/IM \notin S$, then every S-sequence on M in I, can be extended to a maximal one.*

Proof. Suppose that $a_1, a_2, a_3, \ldots$ is a sequence of elements of I such that for any positive integer t, the finite sequence $a_1, \ldots, a_t$ is an S-sequence on M, and look for a contradiction. Then, for any positive integer t, we must have $(a_1, \ldots, a_t) \subset (a_1, \ldots, a_t, a_{t+1})$; otherwise we would have $a_{t+1} \in (a_1, \ldots, a_t)$, so $M/(a_1, \ldots, a_t)M = 0 :_{M/(a_1,\ldots,a_t)M} a_{t+1} \in S$. Hence $M/IM = (M/(a_1, \ldots, a_t)M) \otimes_R R/I \in S$ which is a contradiction. Thus

$$(a_1) \subset (a_1, a_2) \subset \cdots \subset (a_1, \ldots, a_t) \subset (a_1, \ldots, a_t, a_{t+1}) \subset \cdots$$

is an infinite strictly ascending chain of ideals of R, contrary to the fact that R is Noetherian. □

Definition 2.7. An R-module M is said to be zero-divisor module if for any submodule N of M, the set $Z_R(M/N)$ is a finite union of prime ideals in $\operatorname{Ass}_R M/N$.

According to [6, Example 2.2], the class of *ZD*-modules contains finitely generated, Laskerian, weakly Laskerian, linearly compact and Matlis reflexive modules. Also it contains modules whose quotients have finite Goldie dimension and modules with finite support, in particular Artinian modules. Therefore the class of *ZD*-modules is much larger than that of finitely generated modules.

In the following result, we get a relation between S-sequences and local cohomology modules of *ZD*-modules.

Lemma 2.8. *Let $I = (a_1, \ldots, a_k)$ be an ideal of R, S be a Melkersson subcategory with respect to I and M be a ZD-module. Then the following conditions are equivalent:*

(i) $H^i_I(M) \in S$ *for all* $i < t$.

(ii) $\operatorname{Ext}^i_R(R/I, M) \in S$ *for all* $i < t$.

(iii) $\operatorname{Ext}^i_R(N,M) \in S$ *for all* $i < t$ *and each finitely generated module* N *such that* $\operatorname{Supp}_R N \subseteq V(I)$.

(iv) *There is an* S*-sequence on* M *in* I *of length* t.

Proof. By [1, Theorem 2.9], the conditions (i) to (iii) are equivalent. We show by induction that (iv) is equivalent to the other conditions.

Let $t = 1$. Let $a \in I$ be an S-sequence on M. Then $0 :_{\Gamma_I(M)} a = 0 :_M a \in S$, and so $\Gamma_I(M) \in S$ by [1, Theorem 2.3]. Now suppose that $\Gamma_I(M) \in S$. Since $\Gamma_I(M/\Gamma_I(M)) = 0$, it follows by [6, Lemma 2.4] that I contains an element a which is non zero-divisor on $M/\Gamma_I(M)$. The exact sequence $0 \longrightarrow \Gamma_I(M) \longrightarrow M \longrightarrow M/\Gamma_I(M) \longrightarrow 0$ induces the following exact sequence

$$\Gamma_{(a)}(\Gamma_I(M)) \longrightarrow \Gamma_{(a)}(M) \longrightarrow \Gamma_{(a)}(M/\Gamma_I(M)).$$

Because $\Gamma_{(a)}(\Gamma_I(M)) = \Gamma_I(M)$ and $\Gamma_{(a)}(M/\Gamma_I(M)) = 0$, it follows by the above exact sequence that $\Gamma_{(a)}(M) \in S$. Therefore $0 :_M a \in S$, and $a \in I$ is an S-sequence on M.

Now assume inductively that $t > 1$ and the result has been proved for smaller values of t. Let $a_1, \ldots, a_t$ be an S-sequence on M in I. Since $0 :_M a_1 \in S$, it follows by [2, Lemma 2.1] that $\operatorname{Ext}^i_R(N, 0 :_M a_1) \in S$ for all $i \in \mathbb{N}_0$ and any finitely generated R-module N. Therefore by equivalence of (i) and (iii), $H^i_I(0 :_M a_1) \in S$ for all $i < t$. Since $a_2, \ldots, a_t$ is an S-sequence on M/a_1M, it follows by inductive hypothesis that $H^i_I(M/a_1M) \in S$ for all $i < t-1$. Consider the R-homomorphism $f : M \xrightarrow{a_1} M$. Since $H^i_I(\operatorname{Ker} f) \in S$ and $H^{i-1}_I(\operatorname{Coker} f) \in S$ for all $i < t$, it follows by [10, Lemma 3.1] that $(0 :_{H^i_I(M)} a_1) = \operatorname{Ker} H^i_I(f) \in S$ for all $i < t$. Therefore $H^i_I(M) \in S$ for all $i < t$.

Suppose that $H^i_I(M) \in S$ for all $i < t$. Take a_1 as in the proof of the case $t = 1$. Then a_1 is non zero-divisor on $M/\Gamma_I(M)$, and we may replace M with $M/\Gamma_I(M)$ and assume that a_1 is M-regular. The exact sequence $0 \longrightarrow M \xrightarrow{a_1} M \longrightarrow M/a_1M \longrightarrow 0$ induces the following exact sequence

$$\cdots \longrightarrow H^{i-1}_I(M/a_1M) \longrightarrow H^i_I(M) \xrightarrow{a_1} H^i_I(M) \longrightarrow H^i_I(M/a_1M) \longrightarrow \cdots.$$

Hence $H^i_I(M/a_1M) \in S$ for all $i < t-1$. Now it follows by inductive hypothesis that $a_2, \ldots, a_t$ is an S-sequence on M/a_1M. Therefore $a_1, \ldots, a_t$ is an S-sequence on M. □

The following result is a generalization of [1, Lemma 2.14(b)].

Theorem 2.9. *Let S be a Melkersson subcategory with respect to I, and M be a ZD-module such that $M/IM \notin S$. Then the length of all maximal S-sequences on M in I is equal to* $\inf\{i : \operatorname{Ext}^i_R(R/I,M) \notin S\}$.

Proof. In view of 2.6, we suppose that $a_1,\dots,a_t$ is a maximal S-sequence on M in I. If we show that $\operatorname{Ext}^t_R(R/I,M) \notin S$, then the claim follows by 2.8. First we reduce to the case that a_1 is a non zero-divisor element on M. Since $0:_{\Gamma_{(a_1)}(M)} a_1 = 0:_M a_1 \in S$, thus $\Gamma_{(a_1)}(M) \in S$. Hence it follows by [1, Theorem 2.7(b)] that $a_1,\dots,a_t$ is an S-sequence on $M' = M/\Gamma_{(a_1)}(M)$ in I, and a_1 is a regular element on M'. The exact sequence $0 \longrightarrow \Gamma_{(a_1)}(M) \longrightarrow M \longrightarrow M' \longrightarrow 0$ induces the following exact sequence

$$\begin{aligned}\operatorname{Ext}^t_R(R/I,\Gamma_{(a_1)}(M)) \longrightarrow \operatorname{Ext}^t_R(R/I,M) &\longrightarrow \operatorname{Ext}^t_R(R/I,M') \\ &\longrightarrow \operatorname{Ext}^{t+1}_R(R/I,\Gamma_{(a_1)}(M)).\end{aligned}$$

Since $\Gamma_{(a_1)}(M) \in S$, thus $\operatorname{Ext}^i_R(R/I,\Gamma_{(a_1)}(M)) \in S$ for all $i \in \mathbb{N}_0$. Therefore it follows by the above exact sequence that, $\operatorname{Ext}^t_R(R/I,M) \in S$ if and only if $\operatorname{Ext}^t_R(R/I,M') \in S$. Now we may replace M with M' and assume that a_1 is non zero-divisor on M.

We use induction on t. Let $t=1$. We suppose that $\operatorname{Ext}^1_R(R/I,M) \in S$ and look for a contradiction. The exact sequence $0 \longrightarrow M \xrightarrow{a_1} M \longrightarrow M/a_1M \longrightarrow 0$ induces the following exact sequence

$$\begin{aligned}0 \quad\longrightarrow\quad & \operatorname{Hom}_R(R/I,M/a_1M) \longrightarrow \operatorname{Ext}^1_R(R/I,M) \xrightarrow{a_1} \operatorname{Ext}^1_R(R/I,M) \longrightarrow \cdots \\ \longrightarrow\quad & \operatorname{Ext}^{t-1}_R(R/I,M) \longrightarrow \operatorname{Ext}^{t-1}_R(R/I,M/a_1M) \longrightarrow \operatorname{Ext}^t_R(R/I,M).\end{aligned}$$

Thus $\operatorname{Hom}_R(R/I,\Gamma_I(M/a_1M)) = \operatorname{Hom}_R(R/I,M/a_1M) \in S$ and so $\Gamma_I(M/a_1M) \in S$. Since a_1 is a maximal S-sequence on M in I, thus $0:_{M/a_1M} a \notin S$ for all $a \in I$. Hence for any $a \in I$, there is $x_a \in M$ such that $x_a + a_1M \notin S$ and $a(x_a + a_1M) = 0$. Set $X = \{x_a + a_1M : a \in I\}$ and $N = (X)$. Then $N \notin S$ and $I \subseteq Z_R(N) = \bigcup_{\mathfrak{p} \in \operatorname{Ass}_R N} \mathfrak{p}$. So there is $\mathfrak{q} \in \operatorname{Ass}_R N$ such that $I \subseteq \mathfrak{q}$. Thus there is $a \in I$ such that $x_a + a_1M \notin S$ and $I(x_a + a_1M) = 0$. Therefore $x_a + a_1M \in \Gamma_I(M/a_1M)$ and so $\Gamma_I(M/a_1M) \notin S$ which is a contradiction.

Now assume inductively that $t > 1$ and the result has been proved for smaller values of t. Since $a_2,\dots,a_t$ is a maximal S-sequence on M/a_1M in I, it follows by inductive hypothesis that $\operatorname{Ext}_R^{t-1}(R/I, M/a_1M) \notin S$. Also it follows by 2.8 that $\operatorname{Ext}_R^{t-1}(R/I, M) \in S$. Therefore the claim follows by the above exact sequence. □

Definition 2.10. Let S be a Melkersson subcategory with respect to I and M be a ZD-module such that $M/IM \notin S$. The common length of all maximal S-sequences on M in I, is denoted by $S-\operatorname{depth}_I(M)$. If $M/IM \in S$, we set $S-\operatorname{depth}_I(M) = \infty$.

Example 2.11. [1, Example 2.16] Let M be a ZD-module. The following are some examples of $S-\operatorname{depth}_I(M)$ with S as in Example 2.2.

(a) It is the same as ordinary $\operatorname{depth}_I(M)$.
(b) It is the same as $f-\operatorname{depth}_I(M)$ (filter-depth) which has been defined in the local case in [9] and [8, Definition 3.3].
(c) It is the same as $g-\operatorname{depth}_I(M)$ (generalized depth) which has been defined in the local case in [11, Definition 4.2].

The following result is a generalization of [1, Lemma 2.18].

Corollary 2.12. *Let S be a Melkersson subcategory with respect to I, and M be a ZD-module such that $M/IM \notin S$. Then*

$$\begin{aligned} S-\operatorname{depth}_I(M) &= \inf\{i : \operatorname{Ext}_R^i(R/I, M) \notin S\} \\ &= \inf\{i : H_I^i(M) \notin S\}. \end{aligned}$$

Proof. The first equality follows by 2.9. The second equality follows by the first one and 2.8. □

Theorem 2.13. *Let S be the class of Artinian R-modules, or be the class of all R-modules N with $\dim_R N \leq k$, where k is an integer. Let M be a finitely generated R-module or be a ZD-module such that $M/IM \notin S$. Then*

$$\begin{aligned} S-\operatorname{depth}_I(M) &= \inf\{i : \operatorname{Ext}_R^i(R/I, M) \notin S\} \\ &= \inf\{i : H_I^i(M) \notin S\}. \end{aligned}$$

Proof. If M is a ZD-module such that $M/IM \notin S$, then the claim follows by 2.12. So assume that M is a finitely generated R-module such that $M/IM \in S$. It follows by Nakayama's Lemma that

$$\operatorname{Supp}_R \operatorname{Ext}_R^i(R/I, M) \subseteq \operatorname{Supp}_R M \cap V(I) = \operatorname{Supp}_R M/IM.$$

Thus $\operatorname{Ext}^i_R(R/I,M)\in S$ for all $i\in\mathbb{N}_0$. We note that if N is a finitely generated R-module with $\operatorname{Supp}_R N\subseteq \operatorname{Max}(R)$, then N is Artinian; see [13, Lemma 2.1]. Now we have

$$S-\operatorname{depth}_I(M)=\infty=\inf\{i:\operatorname{Ext}^i_R(R/I,M)\notin S\}.$$

By a similar method, one can get the second equality. □

Corollary 2.14. *Let* $\mathfrak{p}\in\operatorname{Spec}(R)$. *Let M be a finitely generated R-module or be a ZD-module such that* $M_{\mathfrak{p}}/\mathfrak{p}M_{\mathfrak{p}}\neq 0$. *Then*

$$\operatorname{depth} M_{\mathfrak{p}}=\inf\{i:\mu^i(\mathfrak{p},M)\neq 0\}.$$

Proof. We note that $\mu^i(\mathfrak{p},M)=\dim_{R_{\mathfrak{p}}}\operatorname{Ext}^i_{R_{\mathfrak{p}}}(R_{\mathfrak{p}}/\mathfrak{p}R_{\mathfrak{p}},M_{\mathfrak{p}})$ for any integer i. □

REFERENCES

[1] M. Aghapournahr, L. Melkersson, *Local cohomology and Serre subcategories*, J. Algebra **320** (2008), 1275-1287.

[2] M. Asgharzadeh, M. Tousi, *A unified approach to local cohomology modules using Serre classes*, Canad. Math. Bull. **53** (2010), 577-586.

[3] M. Brodmann, L. T. Nhan, *A finiteness result for associated primes of certain Ext-modules*, Comm. Algebra **36** (2008), 1527-1536.

[4] M. P. Brodmann, R. Y. Sharp, *Local Cohomology: An Algebraic Introduction with Geometric Applications*, Cambridge University Press, Cambridge, 1998.

[5] W. Bruns, J. Herzog, *Cohen-Macaulay Rings*, Revised Version, Cambridge University Press, Cambridge, 1998.

[6] K. Divaani-Aazar, M. A. Esmkhani, *Artinianness of local cohomology modules of ZD-modules*, Comm. Algebra **33** (2005), 2857-2863.

[7] E. G. Evans, *Zero divisors in Noetherian-like rings*, Trans. Amer. Math. Soc. **155** (1971), 505-512.

[8] R. LÃ¼, Z. Tang, *The f-depth of an ideal on a module*, Proc. Amer. Math. Soc. **130** (2002), 1905-1912.

[9] L. Melkersson, *Some applications of a criterion for Artinianness of a module*, J. Pure Appl. Algebra **101** (1995), 291-303.

[10] L. Melkersson, *Modules cofinite with respect to an ideal*, J. Algebra **285** (2005), 649-668.

[11] L. T. Nhan, *On generalized regular sequences and the finiteness for associated primes of local cohomology modules*, Comm. Algebra **33** (2005), 793-806.

[12] P. Schenzel, N. V. Trung, N. T. Cuong, *Verallgemeinerte Cohen-Macaulay moduln*, Math. Nachr. **85** (1978), 57-73.

[13] Z. Tang, *Local-global principle for the Artinianness of local cohomology modules*, Comm. Algebra **40** (2012), 57-63.

I. K. International University, Postal Code: 34149-1-6818 Qazvin - IRAN
E-mail address: `shpayrovi@ikiu.ac.ir`
E-mail address: `lotfi.parsa@yahoo.com`